

Bladder Management - Intrapartum and Postpartum (including Trial of Void)



- Encourage the woman to void every 2 hours in labour and before beginning active second stage.
- Have a low tolerance for catheterisation if woman unable to void on 2 occasions.
- Catheterise immediately if the bladder is palpable and woman cannot void.
- Do not tape the catheter stretched to the thigh. This will decrease the mobility of the urethra, and decrease the mobility of the balloon in the bladder neck.
- During labour, use only 5-10mL of sterile water in the catheter balloon and ensure it is not the vagina.
- Problems with postpartum voiding are managed using the [Trial of Void Pathways A](#) and [B](#).

1. Purpose

This clinical guideline outlines the requirement for prevention and treatment of peripartum bladder dysfunction at the Women's.

Some degree of voiding dysfunction affects 10-15% of postnatal women and can persist for some time following birth. Five per cent of women have significant and longer lasting dysfunction, which if not recognised in the early peripartum period (birth centre, postnatal ward) may lead to bladder over distension and overflow incontinence resulting in long-term, significant bladder damage and voiding dysfunction.

2. Definitions

Overt bladder retention

This is the inability to pass urine within six hours of birth thus requiring catheterisation, in which volumes greater than normal bladder capacity (normal 400-600mL in females) are drained from the bladder. The woman will often complain of pain and the desire to void, may have overflow incontinence mistaken as stress incontinence or may be asymptomatic particularly if an epidural was used during labour.

Covert bladder retention

The woman is able to void however fails to empty at least 50% of her normal bladder capacity, or a post void residual volume of greater than 150mL. These women will often have frequency and pass volumes of < 100mL.

3. Responsibilities

Staff responsible for intrapartum and postpartum bladder management of a woman should follow this guideline.

4. Guideline

4.1 Risk factors

Women at highest risk include:

- primigravida
- prolonged labour, especially prolonged stage 2
- epidural for labour/birth, irrespective of mode of delivery
- need for catheter in labour
- assisted vaginal birth
- caesarean birth
- perineal injury including haematoma, bruising or tear with inadequate analgesia.

Women without these risk factors may be susceptible to voiding dysfunction; a high index of suspicion must be maintained.

Bladder Management - Intrapartum and Postpartum (including Trial of Void)



4.2 Prevention

Prevention of acute bladder distension

In labour

Encourage the woman to void every 2 hours. Measured volumes are preferred.

If the woman is unable to void on 2 occasions, the threshold for catheterisation is low. If the bladder is palpable and the woman cannot void – catheterise immediately.

A soft catheter is preferable. Do not tape the catheter stretched to the thigh as this will decrease the mobility of the urethra and decrease the mobility of the balloon in the bladder neck.

Inflate the balloon with just 5-10mL of sterile water.

The balloon should be inflated with just 5-10mL of sterile water. If the woman does not have an epidural and catheterisation is merely for the purpose of emptying the bladder prior to a procedure, then an in-out catheter should be considered.

To prevent urethral injury, it is important that the balloon is in the bladder and not merely in the urethra. The taping in of the catheter is of equal importance and must be proximal to the woman's vulva to prevent inadvertent pulling and dislodgement. The Flexi-Trak anchoring device may be used (shown fig 1 below). If unavailable, micropore or similar tape can be used (shown fig 2 below).



Fig 1: Taping using the Flexi-Trak. The blue wings can be trimmed for the woman's comfort - be careful not to trim too short.




Fig 2: Any kind of tape can be used. Anchoring as shown reduces the risk of inadvertent dislodgement. Ensure that drainage is not compromised.

Postpartum

Urine volumes should be representative of patient oral input, keeping in mind normal diuresis in the postpartum woman.

If postpartum urinary retention is suspected or the woman is unable to void, threshold for catheterisation should be low.

Use the Trial of Void sticker (see right) for all postpartum women and manage her voiding ability as advised.

This yellow-bordered sticker is to be placed in the progress notes of all postpartum women and completed by midwives looking after them to clearly document this.

Notes										Sign, Print name Designation	
Delivery date:	/ / @	Mode:	NVB	Vac	Forceps	CS	Weight:				
Perineal Trauma:	N/A	1	2	3A	3B	3C	4	Epis	Swelling		
Post-partum void:	/ / @										
Voided within 6/24 post-delivery?	Yes	No									
Normal urge to void?	Yes	No									
Normal commencement/flow?	Yes	No									
Bladder feels empty post void?	Yes	No									
Voided volume 150mL-700mL	Yes	No									
Signature:	Name:										
IFC removal:	/ / @										
TOV 1:	/ /	Time:	mL								
Normal urge to void?	Yes	No									
Normal commencement/flow?	Yes	No									
Bladder feels empty post void?	Yes	No									
Voided volume 150mL-700mL	Yes	No									
Signature:	Name:										
TOV 2:	/ /	Time:	mL								
Normal urge to void?	Yes	No									
Normal commencement/flow?	Yes	No									
Bladder feels empty post void?	Yes	No									
Voided volume 150mL-700mL	Yes	No									
Signature:	Name:										
If 'No' to any criteria:										<input type="checkbox"/> Commence FBC (input & output) AND <input type="checkbox"/> Refer to Physiotherapy (on-call via switchboard weekends/ public holidays)	
										Signature:	
										Name:	
										Date:	

Bladder Management - Intrapartum and Postpartum (including Trial of Void)



4.3 Diagnosis

A common error is failure to diagnose the bladder distension and incomplete bladder emptying.

Symptoms of voiding dysfunction/retention

Symptoms of voiding dysfunction may include:

- no urge to void
- inability to void within 6 hours of birth or within 6 hours of catheter removal urinary frequency, urgency
- lower abdominal pain
- palpable bladder
- overflow incontinence
- voided volumes of <100mL.
- poor commencement /flow of urine
- sense of incomplete bladder emptying

If a woman experiences the above symptoms, commence a Fluid Balance Assessment Chart (MR2054) and refer to physiotherapy using Internal Referral (OP20), notify the physiotherapist (ext 3160), via pager 53167 or using the on call service available via switchboard (on weekends and public holidays).

In order to assess bladder function, the first postpartum void must be recorded, and the woman asked about urge, flow and feeling of complete emptying.

If an IDC has been inserted, document the planned removal time and ensure that 2 formal trial of voids are completed.

If trial of void criteria are not met, commence a Fluid Balance Assessment Chart (MR2054) and a referral to physiotherapy actioned.

Further management is guided by the [Trial of Void pathways A](#) and [B](#).

4.4 Treatment

Refer to [Trial of Void pathways A](#) and [B](#).

Physiotherapist	Phone: (03) 8345 3160
Continence nurse advisor	Phone: (03) 8345 3144
(urogynaecology outpatient clinic)	

5. Evaluation, monitoring and reporting of compliance to this guideline

Compliance to this guideline or procedure will be monitored, evaluated and reported through clinical incidents reported through VHIMS.

Bladder Management - Intrapartum and Postpartum (including Trial of Void)



the women's
the royal women's hospital

Bladder Management - Intrapartum and Postpartum (including Trial of Void)



the women's
the royal women's hospital

6. References

1. Andolf E. Iosif CS. Jorgensen C. Rydhstrom H. Insidious urinary retention after vaginal delivery: prevalence and symptoms at follow-up in a population-based study. *Gynecologic & Obstetric Investigation*. 38(1):51-3, 1994.
2. Ramsay IN. Torbet TE. Incidence of abnormal voiding parameters in the immediate postpartum period. *Neurourology & Urodynamics*. 12(2):179-83, 1993.
3. Malinowski J. Bladder assessment in the postpartum patient. *JOGN Nursing*. 7(4):14-6, 1978.
4. Groutz A. Gordon D. Wolman I. Jaffa A. Kupferminc MJ. Lessing JB. Persistent postpartum urinary retention in contemporary obstetric practice. Definition, prevalence and clinical implications. *Journal of Reproductive Medicine*. 46(1):44-8, 2001.
5. Lee SN. Lee CP. Tang OS. Wong WM. Postpartum urinary retention. *International Journal of Gynaecology & Obstetrics*. 66(3):287-8, 1999.
6. Yip SK. Hin LY. Chung TK. Effect of the duration of labor on postpartum postvoid residual bladder volume. *Gynecologic & Obstetric Investigation*. 45(3):177-80, 1998.
7. Jeffery TJ. Thyer B. Tsokos N. Taylor JD. Chronic urinary retention postpartum. *Australian & New Zealand Journal of Obstetrics & Gynaecology*. 30(4):364-6, 1990.

7. Legislation/Regulations related to this guideline

Not applicable.

8. Appendices

Appendix 1: [Maternity Trial of Void- Pathway A](#) (initial management)

Appendix 2: [Maternity Trial of Void-Pathway B](#) (subsequent management)

Please ensure that you adhere to the below disclaimer:

PGP Disclaimer Statement

The Royal Women's Hospital Clinical Guidelines present statements of 'Best Practice' based on thorough evaluation of evidence and are intended for health professionals only. For practitioners outside the Women's this material is made available in good faith as a resource for use by health professionals to draw on in developing their own protocols, guided by published medical evidence. In doing so, practitioners should themselves be familiar with the literature and make their own interpretations of it.


Whilst appreciable care has been taken in the preparation of clinical guidelines which appear on this web page, the Royal Women's Hospital provides these as a service only and does not warrant the accuracy of these guidelines. Any representation implied or expressed concerning the efficacy, appropriateness or suitability of any treatment or product is expressly negated

In view of the possibility of human error and / or advances in medical knowledge, the Royal Women's Hospital cannot and does not warrant that the information contained in the guidelines is in every respect accurate or complete. Accordingly, the Royal Women's Hospital will not be held responsible or liable for any errors or omissions that may be found in any of the information at this site.

You are encouraged to consult other sources in order to confirm the information contained in any of the guidelines and, in the event that medical treatment is required, to take professional, expert advice from a legally qualified and appropriately experienced medical practitioner.

NOTE: Care should be taken when printing any clinical guideline from this site. Updates to these guidelines will take place as necessary. It is therefore advised that regular visits to this site will be needed to access the most current version of these guidelines.

Maternity: Trial of Void- Pathway A



Maternity: Trial of Void- Pathway B

